

物質科学セミナー 2019 年1 月16 日 (水) 午前 10 時 30 分開始
3B213 プレゼンテーションルーム

Physical Chemistry with Single Molecules—Hydrogen-Transfer Dynamics Studied by Scanning Probe Microscopy

Dr. Takashi Kumagai

Group leader, Department of Physical Chemistry, Fritz-Haber Institute

H-atom transfer and H-bond rearrangement are involved in many important physical, chemical, and biological processes such as proton conductor, acid–base reactions, and DNA [1]. However, an accurate and quantitative description of H-atom/bonding dynamics remains a very challenging topic and nuclear quantum effects (NQE) and anharmonicity of the potential energy surface are outstanding problems in the field. In order to examine such dynamics, it is necessary to investigate model systems at the single-molecule level because NQE and anharmonic nature are quite susceptible to a local environment of individual molecules and hidden by inhomogeneities of bulk samples. Low-temperature SPM has provided a new opportunity to study hydrogen-atom/bonding dynamics [2]. Single-molecule tautomerization serves as a fascinating model and I will discuss hydrogen-transfer reactions occurring through various external stimuli as well as quantum tunneling [3].

References

- [1] Hydrogen-transfer reaction. J. T. Hynes, J. P. Klinman, H.-H. Limbach, R. L. Schowen, Wiley-VCH, 2007.
- [2] TK, *Prog. Surf. Sci.* 90, 239 (2015).
- [3] TK *et al. Phys. Rev. Lett.* 111, 246101 (2013).; TK *et al. Nature Chem.* 6, 41 (2014).; J. Ladenthin *et al. ACS Nano* 9, 7287 (2015).; H. Böckmann *et al. Nano Lett.* 16, 1034 (2016).; J. Ladenthin *et al. Nature Chem.* 8, 935 (2016).; M. Koch *et al. J. Am. Chem. Soc.* 139, 12681 (2017).; TK, *J. Chem. Phys.* 148, 102330 (2018).; H. Böckmann *et al. Nano Lett.* 18, 152 (2018).

世話人：物質工学域 近藤剛弘 内線5934

共催:エネルギー物質科学研究センター

