

物質工学域セミナー

2024年10月4日 16:00-

筑波大学 工学系学系3B-213号室

“From photoinduced small polarons to cooperative and macroscopic charge-transfer phase transition, monitored by ultrafast X-ray and infrared techniques”

The charge-transfer RbMn [Fe(CN)₆] materials exhibit wide thermal hysteresis around room temperature between Mn^{III}Fe^{II} tetragonal and Mn^{II}Fe^{III} cubic phases. A single laser shot can drive ultrafast photoinduced charge-transfer phase transition at room temperature, which is persisting. Ultrafast studies, based on X-ray and infrared techniques [1-4], provide fundamental mechanistic insight into far-from-equilibrium electronic and structural dynamics, from photoinduced charge-transfer polarons to tetragonal-to-cubic photoinduced phase transition.

[1] M. Hervé, et al., *Nat. Commun.* **2024**, *15*, 267.

[2] G. Privault, et al., *Angew. Chem. I.E.* **2024**.

[3] G. Azzolina, el al., *Angew. Chem. I.E.* **2021**, *60*, 232267.

[4] G. Azzolina, et al., *J. Mater. Chem. C* **2021**, *9*, 6773xxx



Prof. Eric Collet

Institut Universitaire de France

Univ Rennes

CNRS

**Institut de Physique de Rennes
(Rennes, France)**

問い合わせ先

所裕子, Hiroko Tokoro

(tokoro@ims.tsukuba.ac.jp)

羽田真毅, Masaki Hada

(hada.masaki.fm@u.tsukuba.ac.jp)

王駿豪, Junhao Wang

(wang.junhao.gb@u.tsukuba.ac.jp)